

STEFAN CEL MARE UNIVERSITY OF SUCEAVA

**Journal
Food and Environment Safety
of the
Suceava University**

FOOD ENGINEERING

*Volume XVI, Issue 4
30 DECEMBER 2017*

*Available online
<http://www.fia.usv.ro/fiajournal>*

©

**Published by
University Publishing House of Suceava**

AIMS AND SCOPE

The scientific journal *Food and Environment Safety* is a quarterly publication (4 issues per year) of the Faculty of Food Engineering, Stefan cel Mare University of Suceava, Romania, that is indexed in the International Databases: Index Copernicus Journals Master List, Ulrich's and Chemical Abstract (CAS). Also, it is acknowledged B+ by the main Romanian funding organization for university and postgraduate research programmes - National University Research Council (CNCSIS).

The purpose of *Food and Environment Safety* journal is to provide a means of rapid publication for significant contributions to the improvement and diversification of specific activities in the field of food engineering, food and environment safety, biotechnologies, chemical engineering, biosensors, agriculture and applied sciences as well. Moreover, the journal assures promotion of research results in the field of food production and, from the nutritional and toxicological point of view, it underlines the importance of applying the best analyzing and control methods for consumer's protection, food technologies, industrial biotechnologies and environment protection as well to increase life quality.

Food Engineering Faculty journal has been published since 2000 like *Analele Universitatii Stefan cel Mare Suceava, Sectiunea Colegiul Tehnic (ISSN 1583-2295)*. Between 2002 and 2004 the journal was published biannually like *Analele Universitatii Stefan cel Mare Suceava, Sectiunea Inginerie Alimentara (ISSN 1583-2295)*, then from 2005 till 2009, it was issued biannually like *ANNALS of Suceava University - Food Engineering (ISSN 1842-4597)*, indexed in Index Copernicus data base and acknowledged by the National University Research Council (CNCSIS), Romania. In 2010, the Food Engineering Faculty journal changed its name in *Food and Environment Safety (ISSN 2068 - 6609)*.

The cover of the journal was made by **Niculai Moroşan**.

Available online: <http://www.fia.usv.ro/fiajournal>
Address for submission, e-mail fiajournal@fia.usv.ro

CIP description of National Library

The scientific journal *Food and Environment Safety* (ISSN 2068 – 6609) is published by **University Publishing House of Suceava**, Romania.

Mail address of the University Publishing House of Suceava: Stefan cel Mare University, Suceava, Universitatii str. No. 13, 720229, Suceava, Romania

www.usv.ro, <http://www.usv.ro/index.php/ro/1/Editura%20USV/238/3/15>

Tel: +40 230 216 147, **Fax:** +40 0230 520 080

Editorial contact information:

Faculty of Food Engineering, Stefan cel Mare University, Suceava, Universitatii str., No. 13, 720229, Suceava, Romania,

Phone/Fax: +40 230 520267, www.fia.usv.ro

Editor-in-Chief:

Prof. Ph.D,Eng. Sonia AMARIEI (GUTT), gutts@fia.usv.ro

Subscription information

Subscription information can be obtained from:

- **the e-mails:** fiajournal@fia.usv.ro
- **the postal address:**

Faculty of Food Engineering, Stefan cel Mare University - Suceava,
Universitatii str., No. 13, 720229, Suceava, Romania

Copyright © University Publishing House of Suceava. All right reserved

EDITOR-IN-CHIEF:

Sonia AMARIEI (GUTT) Ștefan cel Mare University of Suceava, Romania

EDITORIAL BOARD

Petru ALEXE Dunărea de Jos University of Galați, Romania

Thiery ABRAHAM Enzymes & Derivates, Bruxelles, Belgium

Marcel AVRĂMIUC Ștefan cel Mare University of Suceava, Romania

Șerif BALDIRAN Amasya University, Amasya, Turkey

Elena-Gabriela BAHIRM Dunărea de Jos University of Galați, Romania

Lidija BASA University of Ljubljana, Slovenia

Zoița Mărioara BERINDE Technical University of Cluj Napoca - North University Center of Baia Mare, Romania

Iulian BOBE Tyton BioSciences, Darien, USA

Marija BODROŽA-SOLAROV University Novi Sad, Institute for Food Technology, Serbia

Ioan CHEREJI University of Oradea, Romania

Cristina-Gabriela CONSTANTINESCU Ștefan cel Mare University of Suceava, Romania

Călina Petruța CORNEA University of Agronomic Sciences and Veterinary Medicine in Bucharest, Romania

Toshiki ENOMOTO Ishikawa Prefectural University, Nonoichi, Japan

Dirk FLOTTMAN Technical University of Aalen, Germany

José Antonio GABALDÓN HERNÁNDEZ Catholic University of Murcia (UCAM), Spain

Robert GRAVANI Cornell University, Department of Food Science, USA

Georg GUTT Ștefan cel Mare University of Suceava, Romania

Nicoleta Gabriela HĂDĂRUGĂ Banat University of Agricultural Sciences and Veterinary Medicine Timisoara, Romania

Cristina-Elena HREȚCANU Ștefan cel Mare University of Suceava, Romania

Elena JOSHEVSKA University St. Kliment Ohridski, Bitola, Macedonia

Süleyman KALELI Sakarya University, Sakarya, Turkey

| | |
|----------------------------|--|
| Nijole KAZLAUSKIENE | Institute of Ecology of Nature Research Center, Vilnius, Lithuania |
| Ana LEAHU | Ștefan cel Mare University of Suceava, Romania |
| Anna MARAZ | Corvinus University of Budapest, Hungary |
| Angel MARTINEZ SANMARTIN | Centro Tecnológico Nacional de la Conserva y Alimentación, Murcia, Spain |
| Carmen MORARU | Cornell University, Department of Food Science, USA |
| Ileana Denisa NISTOR | Vasile Alecsandri University of Bacău, Romania |
| Mircea-Adrian OROIAN | Ștefan cel Mare University of Suceava, Romania |
| Mariana-Atena POIANĂ | Banat University of Agricultural Sciences and Veterinary Medicine Timisoara, Romania |
| Regina PROSKUVIENE | Vilnius Pedagogical University, Lithuania |
| Peter RASPOR | University of Ljubljana, Slovenia |
| Gabriela RÂPEANU | Dunărea de Jos University of Galați, Romania |
| Adrian RIVIȘ | Banat University of Agricultural Sciences and Veterinary Medicine Timisoara, Romania |
| Alice ROȘU | Ștefan cel Mare University of Suceava, Romania |
| Sorin SABAU | Tokai University, Sapporo, Japan |
| Alexandru SZÉP | Sapientia — Hungarian University of Transylvania, Miercurea Ciuc, Romania |
| Carmen SOCACIU | University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania |
| Rodica STURZA | Technical University of Moldova, Chisinau, Moldova |
| Gagik TOROSYAN | State Engineering University of Yerevan, Armenia |
| Never TOROSYAN | CSRO Institute, Yerevan, Armenia |
| Yeva TOROSYAN | European Regional Educational Academy, Yerevan, Armenia |
| Felicia TUTULESCU DRAGOMIR | University of Craiova, Romania |
| Igor WINKLER | Yuriy Fedkovych Chernivtsi National University, Ukraine |
| Gheorghe ZGHEREA | Dunărea de Jos University of Galați, Romania |

CONTENTS:

1. **Research on obtaining high β -glucans content from different sources of yeast by harnessing their biologically active potential** 190 - 195
Ionuț AVRĂMIA, Sonia AMARIEI
2. **Quinoa as a substitute for semolina: some aspects and problems of introduction** 196 - 201
Sergiy BORUK, Igor WINKLER, Olga ROMANOVSKA, Olga GERYCH
3. **The interaction between beans hulls and food dye** 202 - 208
Laura Carmen APOSTOL
4. **Enrichment of pasta products using beetroot** 209 - 215
Péter SIPOS, Márta HORVÁTH, Cintia ADÁCSI, Brigitta HORVÁTH, Beáta BABKA, Zoltán GYŐRI
5. **Perspectives of utilization of nano-dispersive materials based on SiO_2 , TIO_2 and SiO_2 - TIO_2 for wine fining** 216 - 221
Anastasiia SACHKO, Igor KOBASA, Olesya MOYSYURA
6. **Determination of dibutyl phthalate (DBP), benzyl butyl phthalate (BBP) and BIS (2-ethylhexyl) phthalate (DEHP) in soft plastic toys and the first survey of the Bulgarian market** 222 - 233
Valentina CHRISTOVA-BAGDASSARIAN, Julieta TISHKOVA, Anton TACHEV
7. **Estimation of heavy metal levels in green leafy vegetables purchased from Suceava** 234 - 238
Ancuța Elena PRISACARU, Laura Carmen APOSTOL, Sorina ROPCIUC
8. **The influence of regular and genetically modified soybeans on postnatal development of rats** 239 - 244
N. N. OMELCHENKO, G. V. DRONIK, I. A. WINKLER, M. S. ROGOZYNSKYI, V. A. KUCHERIAVA
9. **Sensory evaluation of goat milk cheese under different freezing conditions** 245 - 249
Eugenia Mihaela PRICOP
10. **Supplementation of wheat flour with soy flour, sensory and physico-chemical evaluation of fortified biscuits** 250 - 255
Flavia POP, Zorica VOȘGAN, Lucia MIHALESCU
11. **Technology of fresh herbs storage using hydrogel and antioxidant composition** 256 - 261
Olesia PRISS, Viktoria YEVLASH
12. **Research of quality indicators of curd products on basis of protein-herbal clots** 262 - 268
Olena GREK, Alla TYMCHUK, Larisa CHUBENKO, Kira OVSIENKO
13. **Sustainable packaging solutions for organic fresh berries** 269 - 275
Elisabeta Elena TĂNASE, Adina Alexandra BAICU, Vlad Ioan POPA, Amalia Carmen MITELUȚ, Mihaela DRAGHICI, Andreea STAN, Paul Alexandru POPESCU, Mona Elena POPA

- 14. Antioxidant activity and total phenolic content of grape seeds and peels from Romanian varieties** 276 - 281
Mădălina IUGA, Sorina ROPCIUC, Silvia MIRONEASA
- 15. Recyclable porous materials for the uptake of Bisphenol A** 282 – 286
Rodica STURZA, Mihai MELENCIUC, Denisa NISTOR, Farida BOUDISSA, Thiziri TERKANI, Nou-Houda Hadj ABDELKADER, Azzouz ABDELKRIM
- 16. A study into kinetics of biomass blends dewatering as a stage of thermal degradation** 287 - 295
Yaroslav ZASIADKO, Mykola PRYADKO, Pavlo ZASIADKO, Olexiy BULIANDRA
- 17. Author instructions** i - v
- 18. Subscription information** vi



RESEARCH ON THE QUALITY INDICATORS OF CURD PRODUCTS BASED ON PROTEIN-HERBAL CLOTS

Olena GREK¹, *Alla TYMCHUK¹, Larisa CHUBENKO¹, Kira OVSIENKO¹

¹Educational and Research Institute of Food Technology, National University of Food Technologies,
01601, Volodymyrska str.68, Kyiv, Ukraine

*Corresponding author: 589112@ukr.net

Received 29th October 2017, accepted 19th December 2017

Abstract: *The article presents the research of qualitative indicators of curd products with different nutritional ingredients based on protein-herbal clots. The effect of the number of Rumex juice and the duration of thermoacid processing on the process of precipitation of milk proteins was determined. It was established that the introduction of vegetative coagulant in the amount (9 ± 0.5)% at a temperature (93 ... 95) °C and endurance (3 ... 5) min - provides the optimal yield of protein-herbal clot taking into account restrictions according to organoleptic parameters. The effect of white sugar and apple pectin in fiber on the organoleptic, physico-chemical and rheological indicators curd products was investigated. The dietary fibers increase moisture-proof ability and effective viscosity of samples, and white sugar reduces these indexes due to dehydrating properties. The optimal option is to add to the protein-herbal bunch at the same time two components when mixing - white sugar and apple pectin in fiber in quantities of 15% and 2% respectively. Taking into account the influence of individual non-dairy ingredients - Rumex juice of white sugar and apple pectin in fiber on curd products, the performance of the finished product can purposefully be affected.*

Keywords: *protein-herbal clones, sorrel, cheese products, white sugar, apple pectin in cellulose.*

1. Introduction

Extension of assortment and improvement of quality of curd products is carried out as follows main directions:

- improvement of milk-protein concentrates, which are produced without significant changes of technology on existing equipment;
- creation of new types of curd products based on modified dairy raw materials from modern nutritional ingredients;
- use of innovative constructive ideas and technological processes.

Ingredients are generally administered in small quantities, taking into account to their efficacy. Ago special attention should be paid to the issue of preparation for their uniform distribution in raw materials, including milk-protein mixtures. To do

this, you need to evaluate the structural state of raw material or semi-finished. Choose the stage in the process and the modes for making. Take into account the commercial form of the preparation (in the form of a solution, emulsion, dry powder, and granule), concentration (immobilized preparation on the carrier, composition, concentrate) and technological properties (water, fat solubility, salt stability, swelling, degree dispersion) and others [1-2]. Vegetable ingredients are used as a technological component in many production processes. So, the thermoacid coagulation of milk proteins occurs under action, as Serum with acidity (150 ... 200) °T, and coagulants of vegetable origin with relevant characteristics. The latest trend is aimed at expanding the production of milk- protein

products and also modernization of the assortment. Scientists have developed technologies for coagulation of milk proteins as berry raw materials [3-6], and other receptions. They provide a separation process constituents of milk with juice or extracts of plants including wild plants.

Consequently, the aim is to study the qualitative indicators of curd products on the basis of protein-herbal clots with different nutritional ingredients.

The object of the study is the technology of cheese products based on protein-herbal clots and also study of their quality indicators depending on the content of various foods ingredients. The subject of research - protein-herbal clots obtained by thermoacid coagulation of skimmed milk with sorrel juice, food fibers (apple pectin in fiber), white sugar, indicators of quality of model samples of curd products.

The task is:

- to get protein-herbal clots by thermal acid coagulation of fat-free milk in the presence of sorrel juice as a coagulant;
- research on the effect of white sugar and food fibers on organoleptic, physico-chemical and rheological indicators of curd products on the basis of protein-herbal clots.

2. Materials and methods

Sorrel (*Rumex*) was used as a coagulant for protein-herbal clots. It belongs to the genus of one-, two- and perennial herbaceous plants of the family buckwheat, the order of buckwheat flowers [7]. The water content of *Rumex* leaves is (90 ... 92)% protein 1,5%, total carbohydrates (3.0 ... 5.3)%, fiber 1%, minerals, in (mg%): Na (15), K (500), Ca (47), Mg (85), P (90), Fe (2,0). Vitamins, (mg%): C (43 ... 81), B1 (0.19), B2 (0.1), PP (0.3), β -carotene (2.5). The amount of oxalic acid is determined at the level of 360 mg% [8-11]. According to the literature [12, 13] oxalic acid is divided into organic and

inorganic. In the first case, the acid contained in the fresh sorrel is useful for human health substance. It saturates the body with such necessary for the hemorrhage microelements as iron and magnesium. In the inorganic form it is a substance that is formed after heat treatment plants at temperatures above 100 ° C. The acid is coupled with free calcium, forming salts oxalic acid. Oxalates contribute to the formation of stones in the kidneys, the development of articular rheumatism and arthritis, destruction of teeth and bones. With moderate amounts intake of salt of Oxalic acid, it is easily derived from the human body. The harmful amount is 50 mg per 100 g of product [14]. For thermal oxidation deposition from the leaves of *Rumex* received juice in the following sequences. The ground part was sorted, inspected from dirt and mechanical impurities, washed, dried. Chopped to a homogeneous state for (2 ... 3) minutes on the device of the brand DEX DHB-572 power of 750 W.

The raw material where a coagulation process was held is skim milk, which was pasteurized at a temperature (76 ± 2) ° C with a holding (15 ... 20) s. Mass fraction of dry matter is (11.2 ± 0.7)%, protein (3.7 ± 0.2)%, titrated acidity - (17 ± 1.0) ° T, density - 1032 kg / m³.

Previous studies that took into account organoleptic constraints and protein release mass, the optimum amount of vegetative coagulant with pH (3.1 ... 3.5) is determined at the level (9 ± 0.5)% by weight of milk [3, 4]. This amount changes the active acidity of the mixture for providing a balanced isoelectric state of protein in the entire volume. It leads to their active coagulation for classical modes. In heated to a temperature ($93 \dots 95$) ° C, skim milk it was added *Rumex* juice. During the application was moderately stirred and kept (3 ... 5) minutes until a clot was formed. Complex

the effect on high-temperature milk protein and acid reagent leads to the maximum complete their deposition. The process of coagulation was set visually for intense formation protein clot and serum release. The resulting protein-herbal clot had the following qualitative indicators: the mass fraction of moisture at the level $(64 \pm 2)\%$, titrated acidity $(80 \pm 1)^\circ \text{T}$. The organoleptic characteristics of the clot were as follows. Color was light pistachio, not uniform; consistency of the clot was soft, plastic and tight; taste was milk-protein, cheese, without foreign smells, with a slight herbal flavor. At the next stage of the research, model samples of cheese products were made of proteinherbal clots with white sugar and food fibers (FF). The protein base was added into the mixer at a temperature of $(12 \pm 3)^\circ \text{C}$ and mixed with pre-sieved white sugar after partial mixing; prepared FFs were added to the mixture. If necessary, flavoring substances are introduced to disguise the color in the quantities recommended by producer (apple, kiwi - E600 - E699). The average duration of mixing was (5 ... 10) min.

As an FF, apple pectin in fiber was used (APF), which is produced in Ukraine according to TU U 30335750.001-2000. APF contains extruded wheat bran - 60%, apple powder - 38%, pectin - 2%. Chemical composition of 100 g above mentioned the food fibers the following, in%: the mass fraction of moisture – 6.0; Sugar - 48.7, including reducing – 40.9; sucrose - 7.8, organic acids (by apple acid) - 5.0; cellulose- 13.4, pectin substances – 12.4, including water-soluble – 5.1, water-insoluble – 7.3; proteins - 6.7; fats - 6.8; potassium - 0.08; magnesium - 0.73; calcium – 0.32; phosphorus - 0.240. The vitamin complex of APF is represented by carotenoids (0.8 mg) and ascorbic acid (76.4 mg) [15].

During the experiment, the physical and chemical parameters of the model samples

curd products were studied. Mass fraction of moisture was determined by drying the samples to a constant mass on the laboratory electronic scales-moisture series of ADS manufactured by the company "AHIS". Titrated acidity - a titration method based on neutralization of acidic substances contained in 100 g of product with a solution of sodium hydroxide or potassium concentration of $0.1 \text{ mol} / \text{dm}^3$. Moisture-retaining ability (HU) - by Grauhamm's method of modifications A.A. Alekseev, based on the determination of the amount (mass) of water released from the product is lightly compressed and absorbed by a filter paper. Effective viscosity - on a rotating viscometer "Reotest II". The viscometer uses a two-cylinder a measuring system that allows determining the structural mechanical characteristics of the syrup products. The measurement of the shear stress τ (Pa) was carried out with a change in the gradients of the shear voltage γ in the range of $0.33 \dots 145.8 \text{ s}^{-1}$ for direct traffic. The indicators of viscometer data were specified maximum angle of deviation of the arrow on the instrument scale. Displacement voltage (Pa) was calculated by the formula (Eq. 1):

$$\tau = Z \cdot \alpha, \quad (1)$$

where Z is the cylinder constant, $P \cdot \text{unit}$. scale; α is the measure of measurement.

The effective viscosity ($\text{Pa} \cdot \text{c}$) was determined by the formula (Eq. 2):

$$\eta_{\text{ef}} = \frac{\tau}{\gamma}, \quad (2)$$


where γ is the gradient of shear force, c^{-1} [16-18].

All analytical definitions were performed at least three times. Indicator - is average the standard deviation value. The statistical analysis was conducted with the help of Microsoft Excel version 2010.

3. Results and discussion

The influence of the amount of *Rumex* juice on the process of coagulation of proteins in low-fat milk and output of protein-herbal clot has been confirmed.

The output of the latter under the same conditions process and adjusted according to the amount of dry matter prepared *Rumex* is presented on Fig. 1.



* Mass fraction of moisture of protein-grassy clot - $(64 \pm 2) \%$

Fig. 1. The yield of a protein-grassy bunch, depending on the amount of *Rumex* juice

Organoleptic parameters are a limiting factor due to color variations of clots from barely green to intensely green (for 11% juice). Not formed clot that complicates the process of self-pressing (for 5% juice) is also a limiting factor. Number juice at the level $(9 \pm 0.5)\%$, taking into account the above restrictions, is optimal. Entrance Protein-herbal clot is (203 ± 5) g of 1000 g of skimmed milk. As for the course process, the result of coagulation is a partial dehydration, a decrease in the charge hydrated protein molecules and rapid precipitation in the precipitate. Curd products with a polycomponent composition should be elastic and plastic. This will allow the formation and packaging of products commonly accepted in the industry way on existing equipment [19]. As a result of the research, it was discovered the effect of the individual

prescription components of sugar and food fiber with the above water absorption capacity on quality indicators of cheese products on the basis of protein-herbal clots. So, the amount of white sugar in the recipes of the classic soured items the range varies from 10% to 20% (creams, syrup masses, glazed curds, semi-finished products, etc.). To stabilize the moisture and enrichment of protein clots was Used APF. The basis of the distribution of ingredients is the need to clarify the effects of individual components of non-milk origin and their compositions on physico-chemical and rheological indicators of curd products. Results of determining the change in effective viscosity, moisture retention capacity and titrated acidity of the samples under study are presented in Table. 1.

Table 1
Dependency of titratable acidity, moisture-retaining ability and effective viscosity on the composition of curd products based on protein-grassy clots

| № Sample | Ratio of prescription components in curd products, % | | | Indicators of curd products | | |
|----------|--|-------------|-----|-----------------------------|-------------------------------|-----------------------------|
| | Protein-herbal clots | White sugar | APF | Titrated acidity, °T | Moisture-retaining ability, % | Effective viscosity, Pa · s |
| 1 | 100 | - | - | 80 ± 1 | 57.5 ± 1.3 | 353.3 ± 3.3 |
| 2 | 90 | 10 | - | 78 ± 1 | 49.7 ± 1.0 | 293.5 ± 2.8 |
| 3 | 85 | 15 | - | 77 ± 1 | 46.4 ± 1.1 | 264.2 ± 2.7 |
| 4 | 80 | 20 | - | 76 ± 1 | 43.2 ± 1.3 | 234.8 ± 3.0 |
| 6 | 99 | - | 1 | 80 ± 1 | 60.3 ± 1.2 | 372.2 ± 3.0 |
| 7 | 98 | - | 2 | 79 ± 1 | 66.5 ± 1.1 | 411.3 ± 1.9 |
| 8 | 97 | - | 3 | 79 ± 1 | 71.8 ± 1.4 | 469.6 ± 2.5 |
| 10 | 89 | 10 | 1 | 79 ± 1 | 57.3 ± 2.1 | 345.99 ± 2.2 |
| 11 | 83 | 15 | 2 | 78 ± 1 | 61.1 ± 1.7 | 381.60 ± 2.5 |
| 12 | 77 | 20 | 3 | 77 ± 1 | 67.2 ± 1.6 | 401.03 ± 2.8 |

According to experimental data (Table 1), non-dairy ingredients that are added to the protein-herbal clot (namely, white sugar and / or APF), do not amplify the course of processes and do not activate the increase of acidity. This indicator basically depends from the constituent and qualitative parameters of the protein concentrate obtained by the way thermal coagulation of milk proteins with *Rumex* juice. White sugar slightly more reduces the acidity of the mixture than the APF. Samples containing 20% sugar and 80% protein-herbal clots have acidity of 3 ° T less than samples with the same amount protein component and 3% APF. The APF lowers the moisture content of the model samples, since they consist of hollow ones cells whose walls are cellulose. Other components are hemicellulose and pectin, which are in mostly located between empty cells. Hydrophilic substances of food fibers interact with moisture of dairy products due to the formation of H-bound polyasocyanates with the participation of water molecules and H-bound functional

groups of hydrophilic substances. Molecules Hydrophobic groups become more ordered, as evidenced by a decrease in entropy. Due to dispersion forces hydrophobic groups are aggregated. When combined with moisture, they quickly get it adsorb, increase in size, which increases the viscosity of the system [20].

One of the most important physico-chemical indicators of model samples of curd products is moisture-retaining ability, which is associated with effective viscosity. Previous studies [19] found that for curd products, the index of moisture retaining ability (MRA) on levels above (82.0 ± 2)% are sufficient. Growth of the MRA leads to too dense consistency.

It is found that white sugar lowers the MRA and the effective viscosity of the model specimens. With adding a sugar to the protein-herbal clots in the amount of 20% of the MRA is reduced by 14.3%, and effective viscosity - almost 118.5 Pa · s. Intensively increase from (60.3 ± 1.2)% to (71.8 ± 1.4)% moisture-retaining ability curd products APF with high water

absorption capacity and hydrophilicity. At added to the protein-herbaceous clot above the above non-dairy ingredient in an amount from 1 to 3%, MRA is increased by 11.5%. At the same time, the effective viscosity increases from (372.2 ± 3.0) Pa · s to (469.6 ± 2.5) Pa · s. The ability of the FF to absorb water is related to the degree hydrophilicity and the number of biopolymers present in them, the nature of the surface and porosity particles, their sizes. They combine in their composition biopolymers with different kinship to water. Thus, pectin substances and some hemicelluloses are classified as hydrophilic colloid. This causes their ability to keep water. However, a significant increase in moisture retaining ability and effective viscosity of sour cream products are also undesirable because of this complication is transportation and mechanical the formation of products. Concerning the conditions of the introduction, the optimal option is to add at the same time two components - white sugar and APF. So, when added to the protein-herbal clot. 15% of white sugar and 2% of food fibers (sample number 11), moisture-retaining ability rises from $(57.3 \pm 2.1)\%$ to $(61.1 \pm 1.7)\%$ (Table 1), and the effective viscosity increases from (345.99 ± 2.2) Pa · s to (381.60 ± 2.5) Pa · s. The samples under study have organoleptic characteristics, which satisfy the requirements for such products.

4. Conclusions

The possibility of using *Rumex* juice in the amount $(9 \pm 0.5)\%$ as a coagulant of milk proteins has been confirmed. This makes it possible to obtain organoleptically suitable protein-herbal clots of thermoacid coagulation of skimmed milk for regimes: temperature- $(93 \dots 95)^\circ\text{C}$, duration - $(3 \dots 5)$ min.

Adding prescription components of sugar and APF to protein-herbal clots in different

ways affects the physico-chemical and rheological indices of model samples of curd products. It has been confirmed that the moisture-retaining ability and effective viscosity of the samples increase food fibers. The process is due to the hydrophilicity of biopolymers. White sugar has dehydrating properties and reduces the moisture-retaining ability and effective viscosity of the samples. The acidity of curd products based on protein-herbal clots is determined by acidity the protein base is from $(76 \pm 1)^\circ\text{T}$ and up to $(80 \pm 1)^\circ\text{T}$. Taking into account the influence of individual components non-dairy origin of the quantity of juice *Rumex*, white sugar and APF for curd products, you can purposefully influence the performance of the finished product. Its addition will make it possible ensure compliance with regulatory documents and production conditions.

5. References

- [1]. BAL-PRILIPKO L.V., VENGLYUK YU.P., ZALETOK S.P., MOISEENKO A.M., Improving the technology of combined food products using high quality ingredients and dietary supplements, *Food Industry*, 2: 23–26, (2013).
- [2]. BAL-PRILIPKO L.V., Innovative Technologies of Quality and Safe Meat Products, 207 p., (2012).
- [3]. GREK O., OUOPRIICHUK O., PSHENYCHNA T., The rationalization of the parameters of milk proteins' thermo acid coagulation by berry coagulants, *Food and Environment Safety*, 1: 1–7, (2017).
- [4]. SCHETININ M.P, KOLTYUGINA O.V., KOSYINKINA A.A., Cheese product with berry components, *Dairy Industry*, 10: 58, (2011).
- [5]. OSTROUMOV L.A., BOBYLIN V.V., SMIRNOVA I.A., RAFALOVICH S.R., Investigation of the process of thermo-acid coagulation of milk using various coagulants, *Storage and processing of agricultural raw materials*, 7: 26–27, (1998).
- [6]. BYCHKOVA M.V., Technology development of production of thermo-acid cheeses with various coagulants, *Polzunovskiy almanac*, 4: 92–94, (2009).
- [7]. CHERNYKH V.P., Pharmaceutical Encyclopaedia, 1632 p., (2010).

- [8]. EVTEFEEV YU. V., ZYKOVICH S.N., Investigation of the chemical composition and nutritional status of Rumex K-1 sorrel, *Bulletin of the Altai State Agrarian University*, 76–80, (2011).
- [9]. VYSOCHINA G.V., Phenolic compounds in the taxonomy and phylogeny of the buckwheat family: diss. Doc. of Biological Sciences: 03.00.05, Novosibirsk: 410 p., (2002).
- [10]. KIM J.M., LEE Y.M., JANG D.S., KIM J., Constituents of stems of Rumex japonicus with advanced glycation end products (AGEs) and rat lens Aldose Reductase (RLAR) Inhibitory activity, *Agric. Chem. Biotechnol*, 49(1): 24–27, (2006).
- [11]. RAO K.V., SUNITHA C., BANJI D., A study on the nutraceuticals from the genus Rumex, *Hygeia journal for drugs and medicines*, 2: 76–88, (2011).
- [12]. PEREIRA C., BARROS L., CARVALHO A., FERREIRA I., Use of UFLC-PDA for the Analysis of Organic Acids in Thirty-Five Species of Food and Medicinal Plants, *Food Anal. Methods*, 6: 1337–1344, (2013).
- [13]. RAO K., CH S., BANJII D., SANDHYA S., SAIKUMAR P., Pharmacognostic Studies on Rumex vesicarius, *Asian Journal of Plant Science and Research*, 1: 102–115, (2011).
- [14]. VILIGORSKAYA K.V., KUHLINA O.S., Dismetabolic disorders of the exchange of oxalic acid as a provocative factor for the development of chronic obstructive pulmonary disease, *Achievements of clinical and experimental medicine*, 4: 26–28, (2014).
- [15]. PERFILOVA O.V., MITROFIN M.A., Use of powders from fruit and vegetable extracts for the purpose of expansion of assortment of flour confectionery products, *Achievements of science and technology of agrarian and industrial complex*, 8: 48–50, (2008).
- [16]. ONOPRIICHUK O.O., Improving the technology of cheese products with grain ingredients: diss. cand. tech Sciences: 05.18.16. «Food technology», Kiev: 270 p., (2008).
- [17]. KOSOY V. D., Control of quality of dairy products by methods of physico-chemical mechanics, 208 p., (2005).
- [18]. GREK O.V., YUSHCHENKO N.M., OSMAK T.G., ONOPRIICHUK O.O., RYBAK O.M., TYMCHUK A.V., KRASULYA O.O., Workshop on milk and dairy technology: Teaching. Manual, 431 p., (2012).
- [19]. TYMCHUK A.V., ONOPRIICHUK O.O., GREK O.V., PUHLYAK A.G., PASICHNY V.M., Modern approach to the development of cheese products, *Food industry of agroindustrial complex*, 1: 25–29, (2015).
- [20]. GREK, O.V., KRASULYA O.O., Determination of forms of bound moisture in serum drinks with food fibers, *Food industry*, 14: 58–62, (2013).